

CP violation in $B_{d,s} \rightarrow l^+l^-$ in the model III 2HDM

Yuan-Ben Dai, Chao-Shang Huang, Jian-Tao Li, Wen-Jun Li

Institute of Theoretical Physics, Academia Sinica, Beijing 100080, China

(Dated: February 1, 2008)

We have calculated the Wilson coefficients C_{10}, C_{Q_i} ($i=1,2$) in the $\overline{MS}$ renormalization scheme in the model III 2HDM. Using the obtained Wilson coefficients, we have analyzed the CP violation in decays $B_q^0 \rightarrow l^+l^-$ ($q=d,s$) in the model. The CP asymmetry, A_{CP} , depends on the parameters of models and A_{CP} in $B_d \rightarrow l^+l^-$ can be as large as 40% and 35% for $l=\tau$ and $l=\mu$ respectively. It can reach 4% for B_s^0 decays. Because in SM CP violation is smaller than or equal to $O(10^{-3})$ which is unobservably small, an observation of CP asymmetry in the decays $B_q^0 \rightarrow l^+l^-$ ($q=d,s$) would unambiguously signal the existence of new physics.

PACS numbers: 11.30.E, 13.20.H, 12.60.F, 12.60.J

I. INTRODUCTION

The flavor changing neutral current process, $B_{d,s} \rightarrow l^+l^-$ ($l=\mu, \tau$), has attracted a lot of attention since it is very sensitive to the structure of SM and potential new physics beyond SM and was shown to be powerful to shed light on the existence of new physics before possible new particles are produced at colliders [1, 2, 3, 4, 5]. For example, in a very large region of parameter space supersymmetric (SUSY) contributions were shown to be easy to overwhelm the SM contribution [1, 2, 3, 4, 5] and even reach, e.g., for $l=\mu$, the experimental upper bound [6]

$$\begin{aligned} B_r(B_d \rightarrow \mu^+\mu^-) &< 6.8 \times 10^{-7} \quad (CL = 90\%) \\ B_r(B_s \rightarrow \mu^+\mu^-) &< 2.0 \times 10^{-6} \quad (CL = 90\%). \end{aligned} \quad (1)$$

In other words measuring the branching ratio of $B_{d,s} \rightarrow l^+l^-$ can give stringent constraints on the parameter space of new models beyond SM, especially for that of the minimal supersymmetric standard model (MSSM) because of the $\tan^3\beta$ dependence of SUSY contributions in some large $\tan\beta$ regions of the parameter space [1, 3, 4, 7]. In the model II two Higgs doublet model (2HDM), the branching ratio of $B_s \rightarrow \mu^+\mu^-$ can also reach 2×10^{-8} , which is within good reach at Tevatron Run II, if $\tan\beta$ is large enough (say, ≥ 60) [4, 8]. Comparing with hadronic decays of B mesons, this process is very clean and the only nonperturbative quantity involved is the decay constant that can be calculated by using the lattice gauge theory, QCD sum rules etc.

An observable without hadronic uncertainty at all is the CP asymmetry in the process $B_{d,s} \rightarrow l^+l^-$ since the common uncertain decay constant, which is the only source of hadronic uncertainty for the process, cancels out. CP violation in the b -system has been established from measurements of time-dependent asymmetries in $B \rightarrow J/\Psi K$ decays [9, 10]. It is urgent and important to study CP violation in more processes, including the process $B_{d,s} \rightarrow l^+l^-$ which might be measured in the near future. Obviously for the process $B_{d,s} \rightarrow l^+l^-$ there are no direct CP violations since there are no strong phases in the decay amplitude. But it is well known that CP violating effects can survive after taking into account the mixing of the neutral mesons, B^0 and $\bar{B}^0$, in the absence of the strong phases. Recently, it has been shown that CP violation in $B_{d,s} \rightarrow l^+l^-$ ($l=\mu, \tau$) is also an interesting observable for searching for new physics [11, 12]. In the present paper we study it in the model III 2HDM [13, 14]. It is well-known that in the model III 2HDM the couplings involving Higgs bosons and fermions can have complex phases, which can induce CP violation effects, even in the simplest case in which all tree-level FCNC couplings are negligible. So it is expected that a significant CP asymmetry in $B_{d,s} \rightarrow l^+l^-$ ($l=\mu, \tau$) should exist due to the effects of such extra phases in the model. Another motivation to consider the model III is that one starts in the minimal supersymmetric standard model (MSSM) with a Higgs sector of type II 2HDM and, after integration of squarks and gluinos, one ends up with an unconstrained model III 2HDM [15]. The Wilson coefficients of operators relevant to $b \rightarrow sl^+l^-$ in the effective Hamiltonian have been calculated in the model III 2HDM using the on-shell renormalization prescription [16]. Some Feynman diagrams containing the coupling of a neutral Higgs boson to the charged Higgs and Goldstone bosons were missed in the calculations in [16]. We calculate them, including all the contributions, based on the $\overline{MS}$ renormalization scheme. We then analyze the CP asymmetry in $B_{d,s} \rightarrow l^+l^-$ ($l=\mu, \tau$) using the Wilson coefficients obtained by us. It is shown that the CP asymmetry can be as large as 40% for B_d^0 and 4% for B_s^0 in the reasonable region of parameters in the model.

The organization of the paper is as follows. In Section II we describe the model III 2HDM briefly. In section III we give the effective Hamiltonian responsible for $b \rightarrow sl^+l^-$ in the model. We present the formula for CP asymmetry

in $B_{d,s} \rightarrow l^+ l^-$ in Section IV. The Section V is devoted to numerical results. In Section VI we draw conclusions and discussions. Finally we give contributions to Wilson coefficients C_{10}, C_{Q_i} from individual diagrams in the model III 2HDM in the appendix.

II. THE MODEL III TWO HIGGS DOUBLET MODEL

In general one can have a Yukawa Lagrangian of the form

$$\mathcal{L}_Y = \eta_{ij}^U \bar{Q}_{i,L} \tilde{H}_1 U_{j,R} + \eta_{ij}^D \bar{Q}_{i,L} H_1 D_{j,R} + \xi_{ij}^U \bar{Q}_{i,L} \tilde{H}_2 U_{j,R} + \xi_{ij}^D \bar{Q}_{i,L} H_2 D_{j,R} + h.c. \quad (2)$$

where H_i , for $i=1,2$, are the two scalar doublets of a 2HDM, while $\eta^{U,D}$ and $\xi^{U,D}$ are the non-diagonal matrices of the Yukawa couplings.

For convenience we can choose to express H_1 and H_2 in a suitable basis such that only the $\eta_{ij}^{U,D}$ couplings generate the fermion masses, i.e. such that

$$\langle H_1 \rangle = \begin{pmatrix} 0 \\ \frac{v}{\sqrt{2}} \end{pmatrix}, \quad \langle H_2 \rangle = 0. \quad (3)$$

The two doublets in the basis are of the form

$$H_1 = \frac{1}{\sqrt{2}} \left[\begin{pmatrix} 0 \\ v + \phi_1^0 \end{pmatrix} + \begin{pmatrix} \sqrt{2} G^+ \\ i G^0 \end{pmatrix} \right]; \quad H_2 = \frac{1}{\sqrt{2}} \begin{pmatrix} \sqrt{2} H^+ \\ \phi_2^0 + i A^0 \end{pmatrix}. \quad (4)$$

where $G^{0,\pm}$ are the Goldstone bosons, $H^\pm$ and A^0 are the physical charged-Higgs boson and CP-odd neutral Higgs boson respectively. The advantage of using the basis is that the first doublet H_1 corresponds to the scalar doublet of the SM while the new Higgs fields arise from the second doublet H_2 . So ϕ_2^0 do not have couplings to the gauge bosons of the form $\phi_2^0 ZZ$ or $\phi_2^0 W^+ W^-$.

In eq. (4) ϕ_1^0, ϕ_2^0 are not the neutral mass eigenstates but linear combinations of the CP-even neutral Higgs boson mass eigenstates, H^0 and h^0 :

$$\begin{aligned} H^0 &= \phi_1^0 \cos \alpha + \phi_2^0 \sin \alpha \\ h^0 &= -\phi_1^0 \sin \alpha + \phi_2^0 \cos \alpha \end{aligned} \quad (5)$$

where α is the mixing angle, such that for $\alpha=0$, (ϕ_1^0, ϕ_2^0) coincide with the mass eigenstates.

After diagonalizing the mass matrix of the quark fields, the flavor changing (FC) part of the Yukawa Lagrangian becomes

$$\mathcal{L}_{Y,FC} = \hat{\xi}_{ij}^U \bar{Q}_{i,L} \tilde{H}_2 U_{j,R} + \hat{\xi}_{ij}^D \bar{Q}_{i,L} H_2 D_{j,R} + h.c. \quad (6)$$

where $Q_{i,L}$, $U_{j,R}$, and $D_{j,R}$ now denote the quark fields which are in mass eigenstates and

$$\hat{\xi}^{U,D} = (V_L^{U,D})^{-1} \cdot \xi^{U,D} \cdot V_R^{U,D}. \quad (7)$$

In eq. (7) $V_{L,R}^{U,D}$ are the rotation matrices acting on the up- and down-type quarks, with left or right chirality respectively, so that $V_{CKM} = (V_L^U)^\dagger V_L^D$ is usual Cabibbo-Kobayashi-Maskawa (CKM) matrix. Feynman rules of Yukawa couplings follows from eq. (6) and can be found in, e.g., ref. [17, 18]. The flavor changing neutral current (FCNC) couplings are given by the matrices $\hat{\xi}^{U,D}$ and the charged FC couplings are given by

$$\begin{aligned} \hat{\xi}_{\text{charged}}^U &= \hat{\xi}^U \cdot V_{CKM} \\ \hat{\xi}_{\text{charged}}^D &= V_{CKM} \cdot \hat{\xi}^D. \end{aligned} \quad (8)$$

Because the definition of the $\xi_{ij}^{U,D}$ couplings is arbitrary, we can take the rotated couplings as the original ones and shall write $\xi^{U,D}$ instead of $\hat{\xi}^{U,D}$ hereafter. It is worth to note that the peculiar form of charged FC couplings, eq. (8), is of an important distinction from popular model I and II and has significant phenomenological effects different from those in models I and II [19].

In order to proceed we assume the Cheng-Sher ansatz [13]

$$\xi_{ij}^{U,D} = \lambda_{ij} \frac{\sqrt{m_i m_j}}{v} \quad (9)$$

which ensures that the FCNC within the first two generations are naturally suppressed by small quark masses. In the ansatz the residual degree of arbitrariness of the FC couplings is expressed through the λ_{ij} parameters which are of order one and need to be constrained by the available experiments. In the paper we choose $\xi_{ij}^{U,D}$ to be diagonal for the sake of simplicity so that besides Higgs boson masses only λ_{tt} and λ_{bb} in the quark sector and λ_{ll} in the lepton sector are the new parameters which enter into the Wilson coefficients relevant to the rare leptonic B decays.

The Yukawa Lagrangian (2) has no discrete symmetry. In general, the Higgs potential in 2HDM which has no discrete symmetry may be CP conserved or CP violated [14, 20]. In order to concentrate on the effects of FC Yukawa couplings we assume the potential is CP conserved. Therefore, the new source of CP violation is only from the Yukawa couplings. In a general basis Φ_i ($i=1,2$) the general CP conserved potential is given by:

$$\begin{aligned} V(\Phi_1, \Phi_2) = & \lambda_1(\Phi_1^\dagger \Phi_1 - v_1^2)^2 + \lambda_2(\Phi_2^\dagger \Phi_2 - v_2^2)^2 \\ & + \lambda_3[(\Phi_1^\dagger \Phi_1 - v_1^2) + (\Phi_2^\dagger \Phi_2 - v_2^2)]^2 \\ & + \lambda_4[(\Phi_1^\dagger \Phi_1)(\Phi_2^\dagger \Phi_2) - (\Phi_1^\dagger \Phi_2)(\Phi_2^\dagger \Phi_1)] \\ & + \lambda_5[\text{Re}(\Phi_1^\dagger \Phi_2) - v_1 v_2]^2 + \lambda_6[\text{Im}(\Phi_1^\dagger \Phi_2)]^2 \\ & + \lambda_7(\Phi_1^\dagger \Phi_1 - v_1^2)(\text{Re}(\Phi_1^\dagger \Phi_2) - v_1 v_2) \\ & + \lambda_8(\Phi_2^\dagger \Phi_2 - v_2^2)(\text{Re}(\Phi_1^\dagger \Phi_2) - v_1 v_2). \end{aligned} \quad (10)$$

in which all the coupling constants λ_i are real due to hermiticity. The minimum of the potential is at

$$\langle \Phi_1 \rangle = \begin{pmatrix} 0 \\ v_1 \end{pmatrix}, \quad \langle \Phi_2 \rangle = \begin{pmatrix} 0 \\ v_2 \end{pmatrix}, \quad (11)$$

thus breaks $SU(2) \times U(1)$ down to $U(1)_{EM}$. For the sake of simplicity and to decrease the number of the parameters we assume $\lambda_1 = \lambda_2$ and $\lambda_7 = \lambda_8 = 2\lambda_3 + \lambda_5/2$ in the potential (10). Then we have seven parameters altogether in the Higgs sector. One of them, $v = \sqrt{v_1^2 + v_2^2}$, is fixed by the W boson mass, $M_W^2 = \frac{1}{2}g^2 v^2$. The others can be expressed in terms of mixing angles α' and β and Higgs masses $m_{H^\pm}, m_{A^0}, m_{H^0}, m_{h^0}$. From the potential, it is straightforward to derive

$$\begin{aligned} g_{A^0 H^- G^+} &= \frac{g(m_{A^0}^2 - m_{H^\pm}^2)}{2m_W}, \\ g_{H^0 H^- G^+} &= \frac{ig \sin(\beta - \alpha')(m_{H^0}^2 - m_{H^\pm}^2)}{2m_W}, \\ g_{h^0 H^- G^+} &= \frac{-ig \cos(\beta - \alpha')(m_{h^0}^2 - m_{H^\pm}^2)}{2m_W}, \end{aligned} \quad (12)$$

which have the same forms as those in a general model II 2HDM [4, 21].

The specific basis (H_1, H_2) , eq. (4), can be obtained by a unitary transformation

$$H = S\Phi, \quad S = \begin{pmatrix} \cos \beta & \sin \beta \\ -\sin \beta & \cos \beta \end{pmatrix}, \quad (13)$$

where $H = (H_1, H_2)^T$, $\Phi = (\Phi_1, \Phi_2)^T$, and $\tan \beta = v_2/v_1$.

III. EFFECTIVE HAMILTONIAN FOR $b \rightarrow s(d)l^+l^-$

The effective Hamiltonian describing the flavor changing processes $b \rightarrow s(d)l^+l^-$ can be defined as

$$H_{eff} = -\frac{4G_F}{\sqrt{2}}\lambda_t \left(\sum_{i=1}^{10} C_i(\mu) O_i(\mu) + \sum_{i=1}^{10} C_{Q_i}(\mu) Q_i(\mu) \right). \quad (14)$$

where $\lambda_t = V_{tb}V_{ts}^*$, O'_i ($i = 1, \dots, 10$) are the same as those given in the ref.[22, 23], and Q'_i 's come from exchanging neutral Higgs bosons and have been given in refs. [1, 24].

The explicit expressions of relevant operators in the Model III 2HDM for the process $B_q \rightarrow l^+ l^-$ (q=d,s) are

$$\begin{aligned} O_{10} &= \frac{e^2}{16\pi^2} (\bar{q}_L^\alpha \gamma_\mu b_L^\alpha) (\bar{l} \gamma_\mu \gamma_5 l) \\ Q_1 &= \frac{e^2}{16\pi^2} (\bar{q}_L^\alpha b_R^\alpha) (\bar{l} l) \\ Q_2 &= \frac{e^2}{16\pi^2} (\bar{q}_L^\alpha b_R^\alpha) (\bar{l} \gamma_5 l) \end{aligned}$$

We calculate their Wilson coefficients at one loop level in the $\overline{MS}$ renormalization scheme and results are given in Appendix. Before proceeding, a remark is in place. In calculations of the Wilson coefficients C_{Q_i} the neutral Higgs boson h^0 (H^0 , A^0)-penguin diagrams with the charged Higgs and Goldstone bosons in the loop, which is missed in ref. [16], should be included. In the Appendix we write the contributions of these diagrams to C_{Q_i} in terms of the general coupling constants $g_{A^0 H^- G^+}$, $g_{H^0 H^- G^+}$ and $g_{h^0 H^- G^+}$. However, in numerical calculations we shall confine ourself to the specific case of the couplings (12).

The QCD corrections to coefficients C_i and C_{Q_i} can be incorporated in the standard way by using the renormalization group equations. Q_i ($i = 1, \dots, 10$) does not mix with O_{10} so that the evolution of C_{10} remains unchanged and is given by [22]

$$C_{10}(m_b) = C_{10}(m_W). \quad (15)$$

It is obvious that operators O_i ($i = 1, \dots, 10$) and Q_i ($i = 3, \dots, 10$) do not mix into Q_1 and Q_2 and also there is no mixing between Q_1 and Q_2 . Therefore, the evolution of C_{Q_1}, C_{Q_2} is controlled by the anomalous dimensions of Q_1, Q_2 respectively [24], i.e.,

$$\begin{aligned} C_{Q_i}(m_b) &= \eta^{-\gamma_Q/\beta_0} C_{Q_i}(m_W), \\ &= 1.24 C_{Q_i}(m_W), \quad i = 1, 2, \end{aligned} \quad (16)$$

where $\gamma_Q = -4$ [25] is the one loop anomalous dimension of $\bar{q}_L b_R$, $\eta = \frac{\alpha_s(m_b)}{\alpha_s(M_W)} \approx 1.72$, and $\beta_0 = 11 - (2/3)n_f = 23/3$.

IV. CP ASYMMETRY IN $B_{d,s} \rightarrow l^+ l^-$

We need to know what kind of CP violating observables can be defined in the process $B_q \rightarrow l^+ l^-$ for q=d, s. At first, direct CP violation is absent in this process due to absence of strong phases. T-odd projection of polarization is a kind of useful tool to probe the CP violating effects, for example, in $B \rightarrow X_s l^+ l^-$ [26, 27, 28]. However for the process we are discussing here, we have actually only one independent momentum and one independent spin which can be chosen as those of l^- , so no T-odd projections can be defined. Unlike the case generally discussed for hadronic final states, for example, that in Ref. [29], the detected final states of l^+ and l^- of this process in experiments are basically two asymptotic energy-momentum eigenstates which are not CP eigenstates. Considering for instance B^0 decays to $l^+ l^-$ in the rest frame of B^0 , due to the energy-momentum conservation we denote the four-momenta of l^- and l^+ as $p = (E, \vec{p})$ and $\bar{p} = (E, -\vec{p})$. Then the angular momentum conservation tells us that $l_L^+ l_R^-$ and $l_R^+ l_L^-$ final states are forbidden. Hence we are left with a pair of CP conjugated final states, $l_L^+ l_L^-$ and $l_R^+ l_R^-$, and the couple of the corresponding CP conjugated process. Therefore, we may define the time dependent CP asymmetries as [11]

$$A_{CP}^1(t) = \frac{\Gamma(B_{phys}^0(t) \rightarrow l_L^+ l_L^-) - \Gamma(\bar{B}_{phys}^0(t) \rightarrow l_R^+ l_R^-)}{\Gamma(B_{phys}^0(t) \rightarrow l_L^+ l_L^-) + \Gamma(\bar{B}_{phys}^0(t) \rightarrow l_R^+ l_R^-)} \quad (17)$$

$$A_{CP}^2(t) = \frac{\Gamma(B_{phys}^0(t) \rightarrow l_R^+ l_R^-) - \Gamma(\bar{B}_{phys}^0(t) \rightarrow l_L^+ l_L^-)}{\Gamma(B_{phys}^0(t) \rightarrow l_R^+ l_R^-) + \Gamma(\bar{B}_{phys}^0(t) \rightarrow l_L^+ l_L^-)} \quad (18)$$

Two corresponding time integrated CP asymmetries are

$$A_{CP}^i = \frac{\int_0^\infty dt \Gamma(B_{phys}^0(t) \rightarrow f_i) - \int_0^\infty dt \Gamma(\bar{B}_{phys}^0(t) \rightarrow \bar{f}_i)}{\int_0^\infty dt \Gamma(B_{phys}^0(t) \rightarrow f_i) + \int_0^\infty dt \Gamma(\bar{B}_{phys}^0(t) \rightarrow \bar{f}_i)} \quad i = 1, 2 \quad (19)$$

Where $f_{1,2} = l_{L,R}^+ l_{L,R}^-$ with $l_{L(R)}$ being the helicity eigenstate of eigenvalue $-1(+1)$, $\bar{f}$ is the CP conjugated state of f .

The time evolutions of the initial pure B^0 and $\bar{B}^0$ states are given by[30]

$$\begin{aligned} |B_{phys}^0(t)\rangle &= g_+(t)|B^0\rangle + \frac{q}{p}g_-(t)|\bar{B}^0\rangle, \\ |\bar{B}_{phys}^0(t)\rangle &= \frac{p}{q}g_-(t)|B^0\rangle + g_+(t)|\bar{B}^0\rangle. \end{aligned} \quad (20)$$

with $g_{\pm}(t)$ given by

$$\begin{aligned} g_+(t) &= \exp(-\frac{1}{2}\Gamma t - imt)\cos(\frac{\Delta m}{2}t), \\ g_-(t) &= \exp(-\frac{1}{2}\Gamma t - imt)isin(\frac{\Delta m}{2}t) \end{aligned} \quad (21)$$

The absence of strong phases implies

$$|A_f| = |\bar{A}_{\bar{f}}|, \quad |A_{\bar{f}}| = |\bar{A}_f| \quad (22)$$

where $A_f(\bar{A}_{\bar{f}}) = \langle f|\mathcal{H}_{eff}|B^0(\bar{B}^0)\rangle$. And the CPT invariance leads to

$$\frac{\bar{A}_f}{A_f} = \left(\frac{A_{\bar{f}}}{\bar{A}_{\bar{f}}}\right)^*. \quad (23)$$

In the above definition of CP asymmetry we need to separate the final state $l_L^+l_L^-$ from $l_R^+l_R^-$ in order to measure CP asymmetry. For $l=\tau$, the polarization analysis is straightforward. However, detecting tau's is difficult experimentally. For $l=\mu$, in principle one can separate the final state $\mu_L^+\mu_L^-$ from $\mu_R^+\mu_R^-$ by measuring the energy spectra of the electron from muon decay[31]. A μ_L will decay to an energetic e_L , which must go forward to carry the muon spin, and a less energetic pair of neutrino and antineutrino because the electron is always left-handed¹ and the energy-momentum and angular momentum are conserved. Due to the same reason, for μ_R , the relative energies of electron and a pair of neutrino and antineutrino are roughly reversed. Therefore, the energy spectra of the electron from the muon decay is a powerful μ spin analyzer. However, in practice muons never decay in a 4π detector because the lifetime of a muon is long ($c\tau=659$ m). As pointed out in ref.[11], a possible way to measure a polarized muon decay is to build special detectors which can make muons lose its energy but keep polarization so that the polarized muon decays can be measured.

In order to make measurements accessible, as proposed in ref. [12], one can define the CP violating observable as

$$\begin{aligned} A_{CP} &= \frac{D}{S}, \\ D &= \int_0^\infty dt \sum_{i=1,2} \Gamma(B_{phys}^0(t) \rightarrow f_i) \\ &\quad - \int_0^\infty dt \sum_{i=1,2} \Gamma(\bar{B}_{phys}^0(t) \rightarrow \bar{f}_i), \\ S &= \int_0^\infty dt \sum_{i=1,2} \Gamma(B_{phys}^0(t) \rightarrow f_i) \\ &\quad + \int_0^\infty dt \sum_{i=1,2} \Gamma(\bar{B}_{phys}^0(t) \rightarrow \bar{f}_i) \end{aligned} \quad (24)$$

Different from A_{CP}^i ($i=1,2$), it is accessible experimental to measure such defined CP asymmetry A_{CP} . So in the following we shall concentrate on it.

Using the effective Hamiltonian, we obtain by a straightforward calculation ²

$$\frac{\bar{A}_{f_1}}{A_{f_1}} = -\frac{\lambda_t}{\lambda_t^*} \frac{C_{Q1}\sqrt{1-4\hat{m}_l^2} + (C_{Q2} + 2\hat{m}_l C_{10})}{C_{Q1}^*\sqrt{1-4\hat{m}_l^2} - (C_{Q2}^* + 2\hat{m}_l C_{10}^*)}, \quad (25)$$

¹ In the present case it is quite a good approximation to ignore the mass of electron.

² We have neglected the contributions, which is smaller than or equal to 10^{-3} of the leading term, from the penguin diagrams with c and u quarks in the loop. It is true for both B_d and B_s decays[33]. Therefore, although there are weak phases from the c or u quark in the loop, in particular, for B_d , the effect on the decay phase induced by them is negligibly small.

where $\lambda_t = V_{tb}V_{td}^*$ or $V_{tb}V_{ts}^*$, $\hat{m}_l = m_l/m_{B^0}$ and C_i 's are the Wilson coefficients at the m_b scale. Because C_{Q_i} 's are proportional to m_l and C_{10} is independent of m_l it follows from eq. (25) that the CP asymmetry in $B_{d,s} \rightarrow l^+l^-$ is independent of the mass of the lepton in the approximation of neglecting $4\hat{m}_l^2$. That is, it is the same for $l = \text{electron, muon}$.

In SM, one has[32]³

$$\frac{q}{p} = -\frac{M_{12}^*}{|M_{12}|} = -\frac{\lambda_t^*}{\lambda_t}, \quad (26)$$

up to the correction smaller than or equal to order of 10^{-2} , C_{10} is real, $C_{Q_2} = 0$, and C_{Q_1} is negligibly small. So it follows from Eqs.(25), (26) that there is no CP violation in SM.⁴ If one includes the correction smaller than order of 10^{-2} to $|q/p|=1$ ⁵ one will have CP violation of order of 10^{-3} for B_d^0 and 10^{-4} for B_s^0 which are unobservably small.

In the approximation $|\frac{q}{p}| = 1$ the time integrated CP asymmetry is

$$A_{CP} = -\frac{2Im(\xi)X_q}{(1+|\xi|^2)(1+X_q^2)}, \quad q = d, s, \quad (27)$$

where $X_q = \frac{\Delta m_q}{\Gamma}(q = d, s \text{ for } B_d^0 \text{ and } B_s^0 \text{ respectively})$, and ξ is

$$\xi = \frac{C_{Q1}\sqrt{1-4\hat{m}_l^2} + (C_{Q2} + 2\hat{m}_l C_{10})}{C_{Q1}^*\sqrt{1-4\hat{m}_l^2} - (C_{Q2}^* + 2\hat{m}_l C_{10}^*)} \quad (28)$$

As expected, it is nonzero in the presence of CP violating phases. In deriving Eq. (27) we have used eq. (26) which is the result for $B^0-\bar{B}^0$ mixing in the SM. $B^0-\bar{B}^0$ mixing in the model II 2HDM has been examined in ref. [34]. In the model III there are new CP violating phases which might affect the mixing. However, for the values of parameters which we assumed (see below, eq.(30)) the correction to the SM value of q/p is below 20% and consequently we can still use eq. (27) as an approximation. Because X_s is larger than 19.0(90% CL) and X_d is just about 0.76 [6] A_{CP} in B_s decays is much smaller than that in B_d decays, as can be seen from eq. (27).

V. NUMERICAL RESULTS

In numerical calculations, the following values of parameters are assumed:

$$m_{A^0} = 120GeV, \quad m_{h^0} = 115GeV, \quad m_{H^0} = 160GeV, \quad m_{H^\pm} = 200GeV. \quad (29)$$

To study the dependence of A_{CP} on masses of Higgs bosons, we also include below some results obtained with the doubled values of eq. (29) for Higgs boson masses.

The plots of $|C_{Q1}|$ and $|C_{Q2}|$ for τ versus $|\lambda_{bb}|$ for $|\lambda_{\tau\tau}| = 5$ and fixed $|\lambda_{tt}|$ are shown in Fig. 1 and Fig. 2 respectively. One can see from the figures that $|C_{Q_i}|$ increases when $|\lambda_{bb}|$ increases and the increase is faster for a larger $|\lambda_{tt}|$ than that for a small $|\lambda_{tt}|$. Comparing with the model II 2HDM, $|C_{Q_i}|$ can reach the values larger than those in the model II if the value of $|\lambda_{bb}|$, taken to be $\gg 1$, is equal to that of $\tan\beta$ in the model II and the parameter $|\lambda_{tt}|$ is larger than $1/|\lambda_{bb}|$. We calculate the constraint on $|\lambda_{bb}|$ and $|\lambda_{tt}|$ for $|\lambda_{\mu\mu}| = 50$ due to the experimental upper bound of $Br(B_s \rightarrow \mu^+\mu^-)$ and the result is shown in Fig. 3 where the horizontal line represents the upper limit of $|\lambda_{tt}|$ which comes from the experimental constraints of $B-\bar{B}$ mixing, $\Gamma(b \rightarrow s\gamma)$, $\Gamma(b \rightarrow c\tau\bar{\nu}_\tau)$, ρ_0 , R_b and electric dipole moments (EDMs) of the electron and neutron [18]. The region under the curve (and the horizontal line) is allowed by the experimental bound of $Br(B_s \rightarrow \mu^+\mu^-)$. We also calculate the corresponding constraint by including the phases of the two parameters. The result is that the constraint is not sensitive to the phase of λ_{bb} or λ_{tt} , as expected.

³ Note that the phase convention between B^0 and $\bar{B}^0$ is fixed as $\mathcal{CP}|B^0\rangle = -|\bar{B}^0\rangle$ when deriving eqs. (25), (26).

⁴ One can check by combining Eqs. (26) and (25) that all freedoms of phase conventions are cancelled out completely in $\frac{q}{p} \frac{\bar{A}_{f1}}{A_{f1}}$, including the one between B^0 and $\bar{B}^0$.

⁵ According to the box diagram calculation in SM, the deviation of $|q/p|$ from 1 is $\sim 10^{-3}(10^{-5})$ for $B_d(B_s)$ [32]. So 10^{-2} is a conservative estimate.

In numerically calculating A_{CP} , the following values of parameters are assumed:

$$|\lambda_{bb}| = 50, \quad |\lambda_{tt}| = 0.02, \quad (30)$$

$$|\lambda_{\tau\tau}| = 5, 50, \quad |\lambda_{\mu\mu}| = 5, 50, \quad \theta_{\tau\tau} = \pi/4, \quad \theta_{\mu\mu} = \pi/4, \quad (31)$$

and

$$\theta_{tt} + \theta_{bb} = \pi/2, \pi/3 \quad (32)$$

has been set for simplicity. The values of Higgs boson masses and $\lambda_{bb,tt}$, i.e., eqs. (29), (30), and (32) have been set to satisfy the experimental constraints of $B - \bar{B}$ mixing, $\Gamma(b \rightarrow s\gamma)$, $\Gamma(b \rightarrow c\tau\bar{\nu}_\tau)$, ρ_0 , R_b and electric dipole moments (EDMs) of the electron and neutron [18]. The value of $|\lambda_{\tau\tau}|$, eq. (31), has been set to satisfy the constraint from $Z \rightarrow l^+l^-$ which has been analyzed and the upper limit is 56 [35]. The Figs. 4 and 5 are devoted to A_{CP} versus θ_{tt} for $B_d^0 \rightarrow \tau^+\tau^-$ and $B_d^0 \rightarrow \mu^+\mu^-$ respectively. A_{CP} depends on θ_{tt} significantly and can reach 40% and 35% for $l = \tau$ and $l = \mu$ respectively. As for the B_s^0 decays, A_{CP} is much smaller than that in B_d^0 decays, e.g., it can only reach 4% for $B_s^0 \rightarrow \tau^+\tau^-$, as expected. In order to see the effects of different parameters we consider two cases (a) and (b) which correspond $|\lambda_{\tau\tau}|$ (or $|\lambda_{\mu\mu}|$) = 5 and 50 respectively and plot two curves in each case in Figs. 4,5. In Fig. 4 the solid (dotted) one corresponds the set of Higgs boson masses same as eq. (29) (doubled values of those in eq. (29)). One can see from Fig. 4a that the maximum of A_{CP} significantly decreases when the masses of Higgs bosons increase. However, Fig. 4b shows that the change is small when the masses of Higgs bosons are doubled. The reason is that in the case (a) $|C_{Q_i}|$ (i=1,2) for low Higgs masses, eq.(29), is the same order of magnitude as $\hat{m}_\tau C_{10}$ and C_{10} is almost real, when Higgs boson masses increase C_{Q_i} decreases but C_{10} keeps unchanged so that the CP asymmetry decreases, as shown in Fig. 4a. In the case (b), C_{Q_i} is much larger than $\hat{m}_\tau C_{10}$. Therefore, when Higgs boson masses increase although C_{Q_i} is decreased ξ is almost unchanged since the numerator and dominator are almost simultaneously scaled (see, eq. (28)). The small change of A_{CP} is due to the increase of $m_{H^\pm}$. In order to see the effect of the phase $\theta_{tt} + \theta_{bb}$ on A_{CP} we plot two curves in Fig. 5 where the solid (dotted) one corresponds $\theta_{tt} + \theta_{bb} = \pi/2$ ($\pi/3$). One can see from the figure that the curve moves toward the direction opposite to that of the transverse axis when $\theta_{tt} + \theta_{bb}$ decreases.

With the branching ratios

$$\begin{aligned} Br(B_q^0 \rightarrow l^+l^-) &= \frac{G_F^2 \alpha_{EM}^2}{64\pi^3} m_{B_q}^3 \tau_{B_q} f_{B_q}^2 |\lambda_t|^2 \sqrt{1 - 4\hat{m}_l^2} \\ &\times \left[(1 - 4\hat{m}_l^2) C_{Q_1}^2 + (C_{Q_2} + 2\hat{m}_l C_{10})^2 \right], \end{aligned} \quad (33)$$

where τ_{B_q} is the B_q lifetime, we calculate the events N_q needed for observing A_{CP} at 1σ in the areas of parameter space in which A_{CP} and the branching ratios both have large values and all experimental constraints are satisfied. For $l=\mu$, they are order of 10^8 and 10^9 for B_d^0 and B_s^0 respectively. Therefore, 10^{10} (10^{11}) B_d (B_s) per year, which is in the designed range in the future B factories with 10^8 - 10^{12} B hadrons per year [20], is needed in order to observe the CP asymmetry in $B \rightarrow \mu^+\mu^-$ with good accuracy. For $l=\tau$, the events N_q are order of 10^6 and 10^7 for B_d^0 and B_s^0 respectively. Assuming a total of 5×10^8 (10^9) $B_d\bar{B}_d$ ($B_s\bar{B}_s$) decays, one can expect to observe ~ 100 identified $B_q \rightarrow \tau^+\tau^-$ events, permitting a test of the predicted CP asymmetry with good accuracy.

VI. CONCLUSIONS

In summary, we have calculated the Wilson coefficients C_{10}, C_{Q_i} (i=1,2) in the $\overline{MS}$ renormalization scheme in the model III 2HDM. Comparing with the model II 2HDM, $|C_{Q_i}|$ can reach the values larger than those in the model II when the value of $|\lambda_{bb}|$, taken to be $\gg 1$, is equal to that of $\tan\beta$ in the model II and the parameter $|\lambda_{tt}|$ is larger than $1/|\lambda_{bb}|$. It is shown that there is a constraint on λ_{bb} and λ_{tt} due to the experimental limit of $Br(B_s \rightarrow \mu^+\mu^-)$.

We have analyzed the CP violation in decays $B_q^0 \rightarrow l^+l^-$ ($q=d,s$). The CP asymmetry depends on the parameters of models, in particular, the phase θ_{tt} significantly. A_{CP} in $B_d \rightarrow l^+l^-$ can be as large as 40% and 35% for $l = \tau$ and $l = \mu$ respectively. It can reach 4% for B_s^0 decays. Because in SM CP violation is smaller than or equal to $O(10^{-3})$ which is unobservably small, an observation of CP asymmetry in the decays $B_q^0 \rightarrow l^+l^-$ ($q = d, s$) would unambiguously signal the existence of new physics.

Acknowledgments

The work was supported in part by the National Nature Science Foundation of China.

Appendix

By computing the self-energy type, Higgs-penguin and box diagrams, $C_{Q_1}^a$ and $C_{Q_2}^a$ for $l = \tau$ with the superscript denoting the type of a diagram or the contribution of exchanging a specific Higgs boson are extracted out and given below

$$\begin{aligned}
C_{Q_1}^{A^0} = & \frac{x_t m_b m_\tau \text{Im} \lambda_{\tau\tau} i}{4m_{A^0}^2 \sin^2 \theta_W} \left\{ \int_0^1 dx \int_0^{1-x} dy \left[\frac{y_t \lambda_{tt}^* (2-x)}{S_1(y_t, x_t)} + \frac{y_t \lambda_{tt} (1+x)}{S_1(x_t, y_t)} \right. \right. \\
& + |\lambda_{tt}|^2 \lambda_{bb} \left(1 + 2 \ln \frac{S_2(y_t) m_{H^-}^2}{\mu^2} + \frac{y_t (\lambda_{tt} - 2i \text{Im} \lambda_{tt} x)}{S_2(y_t) \lambda_{bb}} - \frac{y_t \lambda_{tt}}{S_2(y_t) \lambda_{tt}^*} \right) \\
& + \lambda_{tt}^* \left(1 + 2 \ln \frac{S_2(x_t) m_W^2}{\mu^2} \right) + \frac{\lambda_{bb}}{x_t} \left(1 + 2 \ln \frac{m_{H^-}^2 - S_1(y_t, x_t)}{x_t \mu^2} \right) \\
& + \frac{2y_t g_{A^0 H^- G^+}}{g m_W} \left(\frac{\lambda_{bb} - \lambda_{tt}^* x}{S_1(y_t, x_t)} + \frac{\lambda_{tt} (1-x)}{S_1(x_t, y_t)} \right) - \frac{2[\lambda_{tt}^* + i \text{Im} \lambda_{tt} (2+x_t)x]}{S_2(x_t)} \Big] \\
& + \lambda_{tt} \lambda_{bb}^2 \left[\ln \frac{\mu^2}{m_{H^-}^2} - \int_0^1 dx \ln S_4(y_t) \right] + \lambda_{bb} \left[\ln \frac{\mu^2}{m_W^2} - \int_0^1 dx \ln S_4(x_t) \right] \Big\} \quad (34)
\end{aligned}$$

$$\begin{aligned}
C_{Q_2}^{A^0} = & \frac{x_t m_b m_\tau \text{Re} \lambda_{\tau\tau}}{4m_{A^0}^2 \sin^2 \theta_W} \left\{ \int_0^1 dx \int_0^{1-x} dy \left[\frac{y_t \lambda_{tt}^* (2-x)}{S_1(y_t, x_t)} + \frac{y_t \lambda_{tt} (1+x)}{S_1(x_t, y_t)} \right. \right. \\
& + |\lambda_{tt}|^2 \lambda_{bb} \left(1 + 2 \ln \frac{S_2(y_t) m_{H^-}^2}{\mu^2} + \frac{y_t (\lambda_{tt} - 2i \text{Im} \lambda_{tt} x)}{S_2(y_t) \lambda_{bb}} - \frac{y_t \lambda_{tt}}{S_2(y_t) \lambda_{tt}^*} \right) \\
& + \lambda_{tt}^* \left(1 + 2 \ln \frac{S_2(x_t) m_W^2}{\mu^2} \right) + \frac{\lambda_{bb}}{x_t} \left(1 + 2 \ln \frac{m_{H^-}^2 - S_1(y_t, x_t)}{x_t \mu^2} \right) \\
& + \frac{2y_t g_{A^0 H^- G^+}}{g m_W} \left(\frac{\lambda_{bb} - \lambda_{tt}^* x}{S_1(y_t, x_t)} + \frac{\lambda_{tt} (1-x)}{S_1(x_t, y_t)} \right) - \frac{2[\lambda_{tt}^* + i \text{Im} \lambda_{tt} (2+x_t)x]}{S_2(x_t)} \Big] \\
& + \lambda_{tt} \lambda_{bb}^2 \left[\ln \frac{\mu^2}{m_{H^-}^2} - \int_0^1 dx \ln S_4(y_t) \right] + \lambda_{bb} \left[\ln \frac{\mu^2}{m_W^2} - \int_0^1 dx \ln S_4(x_t) \right] \Big\} \quad (35)
\end{aligned}$$

$$\begin{aligned}
C_{Q_1}^{H^0} = & \frac{x_t m_b m_\tau (\text{Re} \lambda_{\tau\tau} S_\alpha + C_\alpha)}{4m_{H^0}^2 \sin^2 \theta_W} \left\{ \int_0^1 dx \int_0^{1-x} dy \left[\frac{2(S_\alpha (\lambda_{tt}^* - 2 \text{Re} \lambda_{tt} x) + C_\alpha (1-2x))}{S_2(x_t)} \right. \right. \\
& + \frac{S_\alpha y_t \lambda_{tt}^* (x-2)}{S_1(y_t, x_t)} - \frac{2x_t (\text{Re} \lambda_{tt} S_\alpha + C_\alpha) x}{S_2(x_t)} - \frac{y_t \lambda_{tt} S_\alpha (1+x)}{S_1(x_t, y_t)} + \frac{C_\alpha (4x-3x_t)}{x_t S_3(x_t)} \\
& - (\lambda_{tt}^* S_\alpha + C_\alpha) \left(1 + 2 \ln \frac{m_W^2 S_2(x_t)}{\mu^2} \right) - \frac{S_\alpha \lambda_{bb}}{x_t} \left(1 + 2 \ln \frac{m_{H^-}^2 - S_1(y_t, x_t)}{x_t \mu^2} \right) \\
& - \frac{C_\alpha}{x_t} \left(1 + 2 \ln \frac{m_W^2 S_3(x_t)}{\mu^2} \right) + |\lambda_{tt}|^2 \left(\frac{y_t (S_\alpha (\lambda_{tt} - 2 \text{Re} \lambda_{tt} x) + C_\alpha (1-2x))}{S_2(y_t)} \right. \\
& - \frac{y_t \lambda_{bb} (\lambda_{tt} S_\alpha + C_\alpha)}{\lambda_{tt}^* S_2(y_t)} - \frac{\lambda_{bb} (\lambda_{tt}^* S_\alpha + C_\alpha)}{\lambda_{tt}^*} \left. \left(1 + 2 \ln \frac{m_{H^-}^2 - S_2(y_t)}{\mu^2} \right) \right) \\
& + \frac{2y_t g_{H^0 H^- G^+}}{i g m_W} \left(\frac{\lambda_{bb} - \lambda_{tt}^* x}{S_1(y_t, x_t)} + \frac{\lambda_{tt} (1-x)}{S_1(x_t, y_t)} \right) \Big] \\
& + \lambda_{tt} \lambda_{bb} (\lambda_{bb} S_\alpha + C_\alpha) \left[\int_0^1 dx \ln S_4(y_t) - \ln \frac{\mu^2}{m_{H^-}^2} \right] \\
& + (\lambda_{bb} S_\alpha + C_\alpha) \left[\int_0^1 dx \ln S_4(x_t) - \ln \frac{\mu^2}{m_W^2} \right] \Big\} \quad (36)
\end{aligned}$$

$$C_{Q_2}^{H^0} = \frac{x_t m_b m_\tau \text{Im} \lambda_{\tau\tau} S_\alpha i}{4m_{H^0}^2 \sin^2 \theta_W} \left\{ \int_0^1 dx \int_0^{1-x} dy \left[\frac{2(S_\alpha (\lambda_{tt}^* - 2 \text{Re} \lambda_{tt} x) + C_\alpha (1-2x))}{S_2(x_t)} \right. \right.$$

$$\begin{aligned}
& + \frac{S_\alpha y_t \lambda_{tt}^*(x-2)}{S_1(y_t, x_t)} - \frac{2x_t(Re\lambda_{tt}S_\alpha + C_\alpha)x}{S_2(x_t)} - \frac{y_t \lambda_{tt} S_\alpha(1+x)}{S_1(x_t, y_t)} + \frac{C_\alpha(4x-3x_t)}{x_t S_3(x_t)} \\
& - (\lambda_{tt}^* S_\alpha + C_\alpha) \left(1 + 2 \ln \frac{m_W^2 S_2(x_t)}{\mu^2} \right) - \frac{S_\alpha \lambda_{bb}}{x_t} \left(1 + 2 \ln \frac{m_{H^-}^2 - S_1(y_t, x_t)}{x_t \mu^2} \right) \\
& - \frac{C_\alpha}{x_t} \left(1 + 2 \ln \frac{m_W^2 S_3(x_t)}{\mu^2} \right) + |\lambda_{tt}|^2 \left(\frac{y_t(S_\alpha(\lambda_{tt} - 2Re\lambda_{tt}x) + C_\alpha(1-2x))}{S_2(y_t)} \right. \\
& - \left. \frac{y_t \lambda_{bb}(\lambda_{tt} S_\alpha + C_\alpha)}{\lambda_{tt}^* S_2(y_t)} - \frac{\lambda_{bb}(\lambda_{tt}^* S_\alpha + C_\alpha)}{\lambda_{tt}^*} \left(1 + 2 \ln \frac{m_{H^-}^2 - S_2(y_t)}{\mu^2} \right) \right) \\
& + \frac{2y_t g_{H^0 H^- G^+}}{ig m_W} \left(\frac{\lambda_{bb} - \lambda_{tt}^* x}{S_1(y_t, x_t)} + \frac{\lambda_{tt}(1-x)}{S_1(x_t, y_t)} \right) \\
& + \lambda_{tt} \lambda_{bb} (\lambda_{bb} S_\alpha + C_\alpha) \left[\int_0^1 dx \ln S_4(y_t) - \ln \frac{\mu^2}{m_{H^-}^2} \right] \\
& + (\lambda_{bb} S_\alpha + C_\alpha) \left[\int_0^1 dx \ln S_4(x_t) - \ln \frac{\mu^2}{m_W^2} \right] \} \tag{37}
\end{aligned}$$

$$\begin{aligned}
C_{Q_1}^{h^0} &= \frac{x_t m_b m_\tau (Re\lambda_{\tau\tau} C_\alpha - S_\alpha)}{4m_{h^0}^2 \sin^2 \theta_W} \left\{ \int_0^1 dx \int_0^{1-x} dy \left[\frac{2(C_\alpha(\lambda_{tt}^* - 2Re\lambda_{tt}x) + S_\alpha(2x-1))}{S_2(x_t)} \right. \right. \\
& + \frac{C_\alpha y_t \lambda_{tt}^*(x-2)}{S_1(y_t, x_t)} - \frac{2x_t(Re\lambda_{tt}C_\alpha - S_\alpha)x}{S_2(x_t)} - \frac{y_t \lambda_{tt} C_\alpha(1+x)}{S_1(x_t, y_t)} + \frac{S_\alpha(3x_t-4x)}{x_t S_3(x_t)} \\
& - (\lambda_{tt}^* C_\alpha - S_\alpha) \left(1 + 2 \ln \frac{m_W^2 S_2(x_t)}{\mu^2} \right) - \frac{C_\alpha \lambda_{bb}}{x_t} \left(1 + 2 \ln \frac{m_{H^-}^2 - S_1(y_t, x_t)}{x_t \mu^2} \right) \\
& + \frac{S_\alpha}{x_t} \left(1 + 2 \ln \frac{m_W^2 S_3(x_t)}{\mu^2} \right) + |\lambda_{tt}|^2 \left(\frac{y_t(C_\alpha(\lambda_{tt} - 2Re\lambda_{tt}x) + S_\alpha(2x-1))}{S_2(y_t)} \right. \\
& - \left. \frac{y_t \lambda_{bb}(\lambda_{tt} C_\alpha - S_\alpha)}{\lambda_{tt}^* S_2(y_t)} - \frac{\lambda_{bb}(\lambda_{tt}^* C_\alpha - S_\alpha)}{\lambda_{tt}^*} \left(1 + 2 \ln \frac{m_{H^-}^2 - S_2(y_t)}{\mu^2} \right) \right) \\
& - \left. \frac{2y_t g_{h^0 H^- G^+}}{ig m_W} \left(\frac{\lambda_{tt}^* x - \lambda_{bb}}{S_1(y_t, x_t)} + \frac{\lambda_{tt}(x-1)}{S_1(x_t, y_t)} \right) \right] \\
& + \lambda_{tt} \lambda_{bb} (\lambda_{bb} C_\alpha - S_\alpha) \left[\int_0^1 dx \ln S_4(y_t) - \ln \frac{\mu^2}{m_{H^-}^2} \right] \\
& + (\lambda_{bb} C_\alpha - S_\alpha) \left[\int_0^1 dx \ln S_4(x_t) - \ln \frac{\mu^2}{m_W^2} \right] \} \tag{38}
\end{aligned}$$

$$\begin{aligned}
C_{Q_2}^{h^0} &= \frac{x_t m_b m_\tau Im\lambda_{\tau\tau} C_\alpha i}{4m_{h^0}^2 \sin^2 \theta_W} \left\{ \int_0^1 dx \int_0^{1-x} dy \left[\frac{2(C_\alpha(\lambda_{tt}^* - 2Re\lambda_{tt}x) + S_\alpha(2x-1))}{S_2(x_t)} \right. \right. \\
& + \frac{C_\alpha y_t \lambda_{tt}^*(x-2)}{S_1(y_t, x_t)} - \frac{2x_t(Re\lambda_{tt}C_\alpha - S_\alpha)x}{S_2(x_t)} - \frac{y_t \lambda_{tt} C_\alpha(1+x)}{S_1(x_t, y_t)} + \frac{S_\alpha(3x_t-4x)}{x_t S_3(x_t)} \\
& - (\lambda_{tt}^* C_\alpha - S_\alpha) \left(1 + 2 \ln \frac{m_W^2 S_2(x_t)}{\mu^2} \right) - \frac{C_\alpha \lambda_{bb}}{x_t} \left(1 + 2 \ln \frac{m_{H^-}^2 - S_1(y_t, x_t)}{x_t \mu^2} \right) \\
& + \frac{S_\alpha}{x_t} \left(1 + 2 \ln \frac{m_W^2 S_3(x_t)}{\mu^2} \right) + |\lambda_{tt}|^2 \left(\frac{y_t(C_\alpha(\lambda_{tt} - 2Re\lambda_{tt}x) + S_\alpha(2x-1))}{S_2(y_t)} \right. \\
& - \left. \frac{y_t \lambda_{bb}(\lambda_{tt} C_\alpha - S_\alpha)}{\lambda_{tt}^* S_2(y_t)} - \frac{\lambda_{bb}(\lambda_{tt}^* C_\alpha - S_\alpha)}{\lambda_{tt}^*} \left(1 + 2 \ln \frac{m_{H^-}^2 - S_2(y_t)}{\mu^2} \right) \right) \\
& - \left. \frac{2y_t g_{h^0 H^- G^+}}{ig m_W} \left(\frac{\lambda_{tt}^* x - \lambda_{bb}}{S_1(y_t, x_t)} + \frac{\lambda_{tt}(x-1)}{S_1(x_t, y_t)} \right) \right] \\
& + \lambda_{tt} \lambda_{bb} (\lambda_{bb} C_\alpha - S_\alpha) \left[\int_0^1 dx \ln S_4(y_t) - \ln \frac{\mu^2}{m_{H^-}^2} \right] \\
& + (\lambda_{bb} C_\alpha - S_\alpha) \left[\int_0^1 dx \ln S_4(x_t) - \ln \frac{\mu^2}{m_W^2} \right] \}
\end{aligned}$$

$$+ (\lambda_{bb}C_\alpha - S_\alpha) \left[\int_0^1 dx \ln S_4(x_t) - \ln \frac{\mu^2}{m_W^2} \right] \} \quad (39)$$

$$C_{Q_1}^{box} = \frac{x_t m_b m_\tau \lambda_{bb} \lambda_{\tau\tau}^*}{4m_{H^-}^2 \sin^2 \theta_W} \int_0^1 dx \int_0^{1-x} dy \left[\frac{1}{S_1(y_t, x_t)} \right] \quad (40)$$

$$C_{Q_2}^{box} = \frac{-x_t m_b m_\tau \lambda_{bb} \lambda_{\tau\tau}^*}{4m_{H^-}^2 \sin^2 \theta_W} \int_0^1 dx \int_0^{1-x} dy \left[\frac{1}{S_1(y_t, x_t)} \right] \quad (41)$$

Adding all the contributions to C_{Q_1} and C_{Q_2} given above respectively, we get the Wilson coefficients C_{Q_i} ($i=1,2$). In numerical calculations we use the couplings among the neutral and charged Higgs bosons and charged Goldstone bosons given in eq. (12).

The Wilson coefficient C_{10} is

$$\begin{aligned} C_{10} &= C_{10}^{SM} + C_{10}^{new}, \\ C_{10}^{new} &= \frac{x_t |\lambda_{tt}|^2}{48 \sin^2 \theta_W} \left\{ \int_0^1 dx \int_0^{1-x} dy \left[6(2 \sin^2 \theta_W - 1) \ln \frac{m_{H^-}^2 S_3(y_t)}{\mu^2} \right. \right. \\ &\quad - \frac{2y_t(3 - 4 \sin^2 \theta_W)}{S_2(y_t)} - 8 \sin^2 \theta_W \left(1 + \ln \frac{m_{H^-}^2 S_2(y_t)}{\mu^2} \right) \Big] \\ &\quad \left. - (3 - 2 \sin^2 \theta_W) \left[2 \int_0^1 dx (x - 1) \ln S_4(y_t) + \ln \frac{\mu^2}{m_{H^-}^2} \right] \right\} \end{aligned} \quad (42)$$

In above equations, the definitions of the functions S_i are

$$\begin{aligned} S_1(\rho, \phi) &= \rho\phi(1 - x - y) + x\phi + yp \\ S_2(\rho) &= (1 - x - y) + (x + y)\rho \\ S_3(\rho) &= (1 - x - y)\rho + (x + y) \\ S_4(\rho) &= 1 - x + x\rho, \end{aligned} \quad (43)$$

and

$$x_t = \frac{m_t^2}{m_W^2}, \quad y_t = \frac{m_t^2}{m_{H^\pm}^2}, \quad (44)$$

$$C_\alpha = \cos \alpha, \quad S_\alpha = \sin \alpha, \quad (45)$$

with α being the mixing angle of the CP-even neutral Higgs bosons.

-
- [1] C.-S. Huang and Q.-S. Yan, Phys. Lett. **B442** (1998) 209; C.-S. Huang, W. Liao and Q.-S. Yan, Phys. Rev. **D59** (1999) 011701.
 - [2] S.R. Choudhury and N. Gaur, Phys. Lett. **B451** (1999) 86.
 - [3] K.S. Babu and C. Kolda, Phys. Rev. Lett. **84** (2000) 228.
 - [4] C.-S. Huang et al., Phys. Rev. **D63** (2001) 114021; *ibid.* **64** (2001) 059902(E); P.H. Chankowski, L. Slawianowska, P.R. **D63** (2001) 054012; C. Bobeth et al., Phys. Rev. **D64** (2001) 074014; C. Bobeth et al., Phys. Rev. **D66** (2002) 074021; G. Isidori, A. Retico, JHEP **11** (2001) 001.
 - [5] A. Dedes, H. K. Dreiner and U. Nierste, Phys. Rev. Lett. **87** (2001) 251804; R. Arnowitt et al., Phys. Lett. **B538** (2002) 121-129; Z. Xiong, J. M. Yang, Nucl. Phys. **B628** (2002) 193; S. Baek, P. Ko and W. Y. Song, hep-ph/0205259; H. Dreiner, U. Nierste and P. Richardson, hep-ph/0207026; J. K. Mizukoshi, X. Tata and Y. Wang, hep-ph/0208078. For a recent review, see, e.g., Chao-Shang Huang, hep-ph/0210314, a talk presented at 5th International Conference on Hyperons, Charm and Beauty Hadrons, Vancouver, Canada, June 25-29, 2002.

- [6] PDG, Review of Particle Data 2002.
- [7] C. Hamzaoui, M. Pospelov and M. Toharia, Phys. Rev. **D59** (1999) 095005.
- [8] H.E. Logan and U. Nierste, Nucl.Phys.**B586** (2000) 39-55.
- [9] B. Aubert *et al.* [BABAR Collaboration], Phys. Rev. Lett. **87**, 091801 (2001).
- [10] K. Abe *et al.* [Belle Collaboration], Phys. Rev. Lett. **87**, 091802 (2001).
- [11] C.-S. Huang, Liao W., Phys. Lett. **B525** (2002) 107.
- [12] C.-S. Huang, Liao W., Phys. Lett. **B538** (2002) 301.
- [13] T.P. Cheng and M. Sher, Phys. Rev. **D35** (1987) 3484; **D44** (1991) 1461.
- [14] W.S. Hou, Phys. Lett. **B296** (1992) 179; A. Antaramian, L.J. Hall, and A. Rasin, Phys. Rev. Lett. **69**, 1871 (1992); L.J. Hall and S. Weinberg, Phys. Rev. D **48**, R979 (1993); M.J. Savage, Phys. Lett. **B266**, 135 (1991); L. Wolfenstein and Y.L. Wu, Phys. Rev. Lett. **73** (1994) 2809.
- [15] A. Dobado, M.J. Herrero and D. Temes, Phys.Rev. **D65** (2002) 075023; A. Pilaftsis, C.E.M. Wagner, Nucl.Phys. **B553** (1999) 3; M. Carena, J. Ellis, A. Pilaftsis, C.E.M. Wagner, Nucl.Phys. **B586** (2000) 92; A. Dedes and A. Pilaftsis, hep-ph/0209306.
- [16] T.M. Aliev and M. Savci, Phys.Lett. **B481** (2000) 275-286; E.O. Iltan and G. Turan, Phys.Rev. **D63** (2001) 115007.
- [17] D. Atwood, L. Reina and A. Soni, Phys.Rev. **D55** (1997) 3156.
- [18] D. Bowser-Chao, K. Cheung, and W.-Y. Keung, Phys. Rev. **D59** (1999) 115006.
- [19] Z.J. Xiao, C.S. Li, and K.T. Chao, Phys. Lett. **B473** (2000) 148, Phys. Rev. **D62** (2000) 094008; T.M. Aliev, M. Savci, hep-ph/0003188.
- [20] F.J. Botella and J.P. Silva, Phys. Rev. **D51** (1995) 3870-3875; H. Georgi, Hadronic Jour. 1 (1978) 155.
- [21] J. F. Gunion, H. E. Haber, G. Kane and S. Dawson, *The Higgs Hunter's Guide* (Addison-Wesley, Reading, MA, 1990); errata hep-ph/9302272.
- [22] B. Grinstein, M.J. Savage and M.B. Wise, Nucl. Phys. **B319** (1989) 271.
- [23] G. Buchalla, A.J. Buras and M.E. Lauthenbache, Rev. Mod. Phys. **68** (1996) 1125.
- [24] Y.-B. Dai, C.-S. Huang, and H.-W. Huang, Phys. Lett. **B390** (1997) 257.
- [25] C.-S.Huang, *Commun. Theor. Phys.* **2** (1983) 1265.
- [26] F. Krüger and L.M. Sehgal, Phys. Lett. **B380** (1996) 199.
- [27] C-S Huang and LIAO Wei, Phys. Rev. D **61** (2000) 116002; Phys. Rev. D **62** (2000) 016008.
- [28] C.-S. Huang, Z.-H. Zhu, Phys. Rev. D **61** (2000) 015011; Erratum-ibid. D **61** (2000) 119903.
- [29] M. Gronau, Phys. Rev. Lett. **63**(1989) 1451.
- [30] I.I. Bigi *et al.*, in "CP violation", ed. C. Jarlskog (World Scientific, Singapore, 1989).
- [31] A similar analysis was presented for separating t_L from t_R . See, C.R. Schmidt and M.E. Peskin, Phys. Rev. Lett. **69** (1992) 410.
- [32] Y. Nir, hep-ph/9911321 (IASSNS-HEP-99-96).
- [33] A.J. Buras, in: Probing the Standard Model of Particle Interactions, Edited by F. David and R. Gupta, Elsevier Science, Amsterdam, 1999, hep-ph/9806471.
- [34] A.J. Buras, P.H. Chankowski, J. Rosiek, and L. Slawianowska, Nucl. Phys. **B619** (2001) 434.
- [35] E.O. Iltan, Phys. Rev. **D65** (2002) 036003.




FIG. 1: $|C_{Q_1}|$ versus $|\lambda_{bb}|$.




FIG. 2: $|C_{Q_2}|$ versus $|\lambda_{bb}|$.






FIG. 3: The constraint on $|\lambda_{bb}|$ and $|\lambda_{tt}|$ due to the experimental upper bound of $Br(B_s \rightarrow \mu^+ \mu^-)$.




(a)




(b)

FIG. 4: (a) A_{CP} for $B_d^0 \rightarrow \tau^+\tau^-$ versus the CP violating phase θ_{tt} , for $|\lambda_{\tau\tau}| = 5$. The solid line stands for the masses of Higgs bosons same as eq. (29), the dotted line stands for the masses of Higgs bosons same as doubled those in eq. (29). (b) The same as (a), except for $|\lambda_{\tau\tau}| = 50$.



(a)



(b)

FIG. 5: (a) A_{CP} for $B_d^0 \rightarrow \mu^+ \mu^-$ versus the CP violating phase θ_{tt} , for $|\lambda_{\mu\mu}| = 5$. The solid line stands for $\theta_{tt} + \theta_{bb} = \pi/2$, the dotted line for $\theta_{tt} + \theta_{bb} = \pi/3$. (b) The same as (a) except for $|\lambda_{\mu\mu}| = 50$.